

WILDBERRY BLISS BODY LOTION



PHASE	TRADE NAME	INCI	FUNCTION	%(w/w)
A	Water	Water	Solvent	75.20
	Glycerin	Glycerin	Humectant	5.00
	Aristoflex™ Eco T	Caesalpinia Spinosa Gum/Ammonium AMPS Crosspolymer	Rheology Modifier	0.30
	Aristoflex HMB	Ammonium Acryloyldimethyltaurate/ Beheneth-25 Methylacrylate Crosspolymer	Rheology Modifier	0.50
B	Plantasens™ Emulsifier HP10	Sucrose Polysteaate (and) Cetearyl Alcohol (and) Olea Eurpaea Oil Unsaponifiabiles	Emulsification	2.00
	Plantasens™ Olive LD	Hydrogenated Ethylhexyl Olivatate and Hydrogenated Olive Oil Unsaponifiabiles	Emollient	3.00
	Plantasens Cocoa Butter	Theobroma Cacao (Cocoa) Seed Butter	Sensory Additive	5.00
	Beraca Murumuru Butter	Astroceyum murumuru seed butter	Sensory Additive	2.00
	Plantasens VP 170	Ricinus Communis (Casotr) seed oil (and) hydrogenated castor oil (and) copernicia cerifera (Carnauba) wax (and) Cera alba (Beeswax)	Emollient	2.00
	Plantasens Olive Squalane	Squalane	Emollient	2.00
C	Waterin Plus	Saccharide Isomerate (and) Water (and) Glycerin (and) Sodium Levulinate (and) p-Anisic Acid	Active Ingredient	1.00
	Nipaguard™ SCE VITA	Sorbitan Caprylate (and) Propanediol (and) Benzoic Acid	Preservative	1.00
	Wildberry Fragrance Oil	Wildberry Fragrance Oil	Fragrance	1.00

PROCEDURE

1. Combine water and glycerin, begin heating while mixing to 75C.
2. Add RM3 first and mix until fully hydrated.
3. Add RM4 to II and mix until fully hydrated.
4. In separate vessel, combine phase B and heat until 75C.
5. Add IV to III and mix under high stirring for 5 minutes, then begin to cool.
6. Once below 30C, add phase C and homogenize.
7. Adjust pH to 5.5.
8. Take out 1/3 of batch and mix in ____% of [Color solution].